

SM221 – Sample Test #2– Fall 2004

Part 1: Multiple Choice (50%). For each question, circle the letter for the best answer.

1. Let $f(t, p)$ be the speed of sound in meters per second (*mps*) when the temperature is t degrees Celsius ($^{\circ}\text{C}$) and the pressure is p atmospheres (*atm*). The statement

$$\frac{\partial f}{\partial p}(2,10) = 3 \text{ means:}$$

- (a) At a temperature of 2°C and a pressure of 10 atm , the speed of sound is 3 mps .
 - (b) At a temperature of 2°C and a pressure of 10 atm , the speed of sound is increasing with increasing pressure at a rate of 3 mps per atm .
 - (c) At a temperature of 2°C and a pressure of 10 atm , the pressure is at a rate of 3 atm per hour.
 - (d) The speed of sound increases 3 mps for each $^{\circ}\text{C}$ increase in temperature.
 - (e) If the speed of sound increases at 3 mps , the temperature increases at a rate of 2°C per hour and the pressure increase at a rate of 10 atm per hour.
-

2. Suppose the budget b in dollars of a manufacturer is a function of the number T of trucks and number C of cars produces so $b = b(T, C)$. T and C are functions of time t , measure in years, Suppose also that:

$$T(10) = 1000, \quad C(10) = 2000, \quad \frac{dT}{dt}(10) = 500, \quad \frac{dC}{dt}(10) = -400,$$

and:

$$\frac{\partial b}{\partial T}(1000, 2000) = 1000, \quad \frac{\partial b}{\partial C}(1000, 2000) = 800.$$

That rate at which b is increasing with time when $t=10$, in dollars per year, is

- (a) 18000
 - (b) 1800
 - (c) 100
 - (d) 0
 - (e) 180,000
-

3. If $f(x, y) = x^2 + xy$, in which direction is f increasing the fastest at $(3, 1)$?

- (a) $7\vec{i} + 3\vec{j}$
 - (b) $\vec{i} + \vec{j}$
 - (c) $7\vec{i} - 3\vec{j}$
 - (d) $3\vec{i} - 7\vec{j}$
 - (e) equally fast in all directions
-

~~4. Which of the following is a parameterization of the cylinder $x^2 + y^2 = 16$?~~

- ~~(a) $r = 4$~~
 - ~~(b) $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$~~
 - ~~(c) $\vec{r} = 4\cos\theta\vec{i} + 4\sin\theta\vec{j} + z\vec{k}$~~
 - ~~(d) $\vec{r} = x\vec{i} + y\vec{j} + (x^2 + y^2)\vec{k}$~~
 - ~~(e) $r^2 = x^2 + y^2$~~
-

5. The contour map for the function $f(x, y) = \sqrt{x^2 + y^2}$ is a family of :

- (a) paraboloids
 - (b) cones
 - (c) circles
 - (d) right triangles
 - (e) spheres
-

6. An equation of the tangent plane to the surface $z = e^{2x} + \cos(y)$ at the point $(0,0,2)$ is:

- (a) $z=2x+2$ (b) $x+y+z=1$ (c) $z=3x+2y$ (d) $x-y+z=0$ (e) $z=y$
-

7. If $f(x,y) = x^2y - y^2$, then $f_{xy}(2,3)$ equals,

- (a) 0 (b) 2 (c) 3 (d) 4 (e) 6
-

~~8. The tangent plane to the ellipsoid $x^2 + 2y^2 + 3z^2 = 36$ at the point $(1, -2, 3)$ is:~~

- ~~(a) $x - 4y + 9z = 0$ (b) $x + 4y + 9z = 20$ (c) $x - 4y + 9z = 20$ (d) $x - 4y + 9z = 36$ (e) $x + 4y + 9z = 108$~~
-

9. Find a vector in the direction of greatest increase of the function $f(x,y,z) = x^2 - 2xy + z^2$ at the point $(1,1,2)$.

- (a) $\langle 4, -2, 2 \rangle$ (b) $\langle 4, 2, 2 \rangle$ (c) $\langle 2, -2, 4 \rangle$ (d) $\langle 0, -2, 4 \rangle$ (e) $\langle 1, 1, 2 \rangle$
-

10. If $\nabla f(x_0, y_0) = \langle 3, -2 \rangle$, then the direction derivative of f , $D_{\vec{v}}(x_0, y_0)$ in the direction of vector $\vec{v} = \langle 4, 3 \rangle$ is:

- (a) -6 (b) $-\frac{6}{5}$ (c) 0 (d) $\frac{6}{5}$ (e) 6
-

11. If $f = f(x,y)$ is a differentiable function with partial derivatives $f_x(1,2) = 5$ and $f_y(1,2) = -2$, then the directional derivative of f at $(1,2)$ in the direction of the vector $-3\vec{i} + 4\vec{j}$ is:

- (a) -23 (b) $-15\vec{i} + -8\vec{j}$ (c) -23/5 (d) $-3\vec{i} - \frac{8}{5}\vec{j}$ (e) $6\vec{i} + 20\vec{j}$
-

12. The maximum rate of change of the function $f(x, y) = x^2 y$ at the point $(1, 2, 2)$ is

- (a) 3 (b) 5 (c) 4 (d) $\sqrt{5}$ (e) $\sqrt{17}$
-

13. The length of the curve $\langle 2t, \sin(t), \cos(t) \rangle$ for $0 \leq t \leq \pi$ is closest to:

- (a) 2.24 (b) 3.14 (c) 6.59 (d) 7.02 (e) 10.63
-

14. An equation for the plane tangent to the surface $z = e^{2x+y}$ at the point $(0, 0, 1)$ is:

- (a) $z=1$ (b) $z = 2xe^{2x+y} + ye^{2x+y} + 1$ (c) $z=x+y+1$ (d) $z=x+2y+1$ (e) $z=2x+y+1$
-

15. $f(x, y, z) = x^2 y + yz$. The directional derivative of f at the point $(3, -2, 4)$ in the direction of

the unit vector $\left\langle \frac{2}{3}, \frac{2}{3}, -\frac{1}{3} \right\rangle$ is

- (a) -26 (b) -2/3 (c) 0 (d) 4/3 (e) 46/9
-

Part 2: Free Response (50 %). The remaining problems are not multiple choice. Answer them in the space below the problem. Show the details of your work and clearly indicate your answers.

~~16. Find the equation for the plane tangent to the surface $x = u^2 + v^2$, $y = v$, $z = u^2 - v^2$ at the point where $u = 2$ and $v = 1$.~~

17. The function f is differentiable at $(1,2)$. \vec{u} is the unit vector $\left\langle \frac{3}{5}, \frac{4}{5} \right\rangle$ and \vec{v} is the unit vector $\left\langle \frac{5}{13}, \frac{12}{13} \right\rangle$. The direction derivative $D_{\vec{u}}f(1,2) = 1$ and $D_{\vec{v}}f(1,2) = -1$. Compute the gradient vector $\nabla f(1,2)$.

18. Monthly production P in a coal mine is given by $P(K,L) = 8K^{1/2}L^{1/2}$, where P is measured in tons, and K in thousands of dollars spent on equipment, and L in thousands of dollars spent on labor per month. Currently $K=100$ and $L=25$.

- Compute the current production $P(100,25)$.
 - A computation shows the $P_L(100,25) = 8$. In words this means: "For every \$1000 more spent on ...". Complete the sentence carefully.
 - Compute the partial derivative $P_K(100,25)$.
 - Suppose that the amounts spent on equipment and labor are increasing by \$4000 and \$3000 a month, respectively. At what rate is production increasing.
-

19. A charged particle moving in the plane receives acceleration due to an alternating electric field $\vec{a}(t) = 50 \sin(\pi t / 60) \vec{j}$ m/sec². If the initial velocity is $\vec{v}(0) = 100 \vec{i}$ m/sec and the initial position is $\vec{r}(0) = 0$ m,

- Find the position $\vec{r}(t)$.
 - Determine the height at which the particle hits a screen 20 meters down the x axis.
-

20. If $z = x^2y$, $x = t^2 + 1$, $y = t + 1$, use the chain rule to find $\left. \frac{dz}{dt} \right|_{t=2}$

21. The table on the right shows the actual normal take off distance, D , of an A6 Intruder which weighs 35000 lbs with a head wind of 20 knots as a function of temperature, T , in degrees F and altitude, A , in feet.

Temp/Alt	0	2000	4000	6000
0	940	1110	1350	1580
30	1070	1280	1530	1830
60	1280	1510	1810	2140
90	1520	1800	2190	2600

- Estimate $\frac{\partial D}{\partial T}$ when the temperature is 30° and the altitude is 2000 ft.
- Estimate $\frac{\partial D}{\partial A}$ when the temperature is 30° and the altitude is 2000 ft.
- Use these estimates to determine the take off distance if the temperature is 35° and the altitude is 2500 ft.

22. For the following function find the critical point(s) and determine if they are a local maximum, local minimum or saddle points: $f(x,y) = x^3 - 12xy + 8y^3$.

23. For the following function find the critical point(s) and determine if they are a local maximum, local minimum or saddle points: $f(x,y) = 2x^3 + xy^2 + 5x^2 + y^2$.

